

SOLDIER EXPOSURE TO DEPLETED URANIUM AND HUMAN HEALTH RISK ASSESSMENT METHODOLOGY UPDATE - 2010

DAVID P. ALBERTH
SENIOR HEALTH PHYSICIST AND MASTER CONSULTANT

FRANCES SZROM, CHP

HEALTH PHYSICIST AND MASTER CONSULTANT

US ARMY INSTITUTE OF PUBLIC HEALTH
DEPLETED URANIUM (DU) HEALTH EFFECTS AND POLICY WORKSHOP

AFRRI, Bethesda, Maryland
4 November 2010



UNCLASSIFIED

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comments arters Services, Directorate for Info	regarding this burden estimate rmation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington			
1. REPORT DATE 04 NOV 2010	2 DEPORT TYPE			3. DATES COVERED 00-00-2010 to 00-00-2010				
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER			
Soldier Exposure to Depleted Uranium and Human Health Risk					5b. GRANT NUMBER			
Assessment Methodology Update - 2010					5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)					5d. PROJECT NUMBER			
					5e. TASK NUMBER			
				5f. WORK UNIT NUMBER				
U.S. Army Public l	ZATION NAME(S) AND AE Health Command (F Ground,MD,21010	Provisional),5158 Bl	ackhawk Road	8. PERFORMING REPORT NUMB	G ORGANIZATION ER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)				
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)				
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	on unlimited						
13. SUPPLEMENTARY NO	OTES							
14. ABSTRACT								
15. SUBJECT TERMS								
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON				
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	30				


Report Documentation Page

Form Approved OMB No. 0704-0188





INTRODUCTION



- Assessment of depleted uranium (DU) use on the battlefield and potential adverse health effects is a sentinel event in addressing all battlefield environmental and occupational hazards
- Experience with DU to be used for developing policy and procedures for assessing other environmental chemical and embedded metal fragment injuries from combat






BACKGROUND (I)

- Operation Desert Storm First combat use of DU munitions by U.S. Military (1991)
 - Walter Reed Army Medical Center First patient identified with embedded DU metal fragment
 - OTSG Consultants identified through medical records review first 35 patients with fragments from fratricide ("friendly-fire") incidents
 - Initial urine bioassay program to identify patients with undiagnosed, embedded metal fragments





OSAGWI Vision (1998), for DU Issue







BACKGROUND (II)

- MOA between Army OTSG and VA for DU Medical Surveillance and Follow-UP Program (1993)
- Interpretation of laboratory results develops into formal Health Risk Assessment (HRA) Program
- HRA is bridge between lab results and clinical guidance for healthcare provider
- Archiving of letter reports for potential future use in patient assessments
- Determination of what is best for Soldier-patient with potential DU exposure





APPLICATION TO AN INTEGRATED ARMY HEAVY METALS PROGRAM

Experience gained in the DU
Health Risk Assessment Process
is being used as a model for
addressing issues with other
heavy metals





- Assessment of retained metal fragments in Servicemembers and Veterans
- Awareness Training and Effective Risk Communication for Servicemembers, their Families, and their Healthcare Providers
- Integration of AMEDD's Health Hazard Assessment and Human Health Risk Assessment Programs – Requirement for Communication with Munitions Developers





POTENTIAL DU EXPOSURES FROM MILITARY OPERATIONS



- Hard target impact and perforation of armored vehicles by DU munitions
- Vehicle fires involving DU munitions
- Potential for subsequent environmental exposures through resuspension of DU residues







DU EXPOSURE CATEGORIES



I	Soldiers in, on, or near armored vehicle struck and penetrated with DU munitions
II	Soldiers whose MOS requires entering DU-damaged vehicles
Ш	Soldiers with incidental DU exposure










FACTORS AFFECTING DU TOXICITY





- Route of exposure
- Chemical form
- Amount internalized
- Solubility
- Particle size and distribution





ARMY/BATTELLE CAPSTONE DU HUMAN HEALTH RISK ASSESSMENT



FINDINGS

- Greatest potential for inhalation exposure - in, on or near an armored vehicle at time of impact by a DU penetrator (DU Exposure Level I)
- Chemical toxicity may be the more limiting factor for soluble DU forms; however, military DU sources limit the availability of the more soluble forms
- All other potentially exposed individuals received significantly less intake through inhalation (DU Exposure Levels II and III)

See http://fhp.osd.mil/du/du_capstone/index.pdf





NAS COMMITTEE ON TOXICOLOGY PEER REVIEW



"Review of Toxicologic and Radiologic Risks to Military Personnel from Exposure to Depleted Uranium During and After Combat" (2008)

- Army DU Research-Integrated Process Team recommended an independent scientific peer review by the National Academy of Sciences Committee on Toxicology
- See link to the NAS-COT review of the Capstone Reports: http://www.nap.edu/catalog.php?r ecord_id=11979
- Findings and recommendations presented in the Academy's report compared and contrasted with the "Depleted Uranium Aerosols Doses and Risk: Summary of U.S. Assessments" (2005) for close-out of project





THE CAPSTONE DEPLETED URANIUM (DU) AEROSOLS STUDY AND HUMAN **HEALTH RISK ASSESSMENT**

- Final two Data Quality Objectives for this project completed in the year 2009:
 - Review of the Capstone DU Project reports by the NRC, Council, National Academy of Sciences Committee on Toxicology
 - Publication of a series of 14 technical articles on the project in a dedicated, special issue of *Health* Physics, an independent, peer-reviewed scientific journal (Health Physics Journal, The Radiation Safety Journal, Special Issue: The Capstone Depleted Uranium Aerosol Characterization and Risk Assessment Study, ISSN 0017-9078, Vol. 96, No.

	100 may 200 miles (100 miles 100 mil		PHYSICS Safety Journal			
March 2009	Volume 96		Number 3 ISSN 0017-	907		
PART I: THE CAPSTO	NE DEPLETED URANIC ASSE ONE DEPLETED	JM A	ENTS EROSOL CHARACTERIZATION AND RISK ENT STUDY PART II: THE CAPSTONE HUMAN HEAL	ТН		
URANIUM AEROSOL CHARACTERIZATION Overview of the Capatone Depleted Uranium Study of Aerosols from Impact with Armored Vehicles: Test Setup and Aerosol Generation, Characterization, and Application in Assessing Dose and Risk			RISK ASSESSMENT Methods Used to Calculate Doses Resulting from Inhalation of Capstone Depleted Uranium Aerosols Acrosols Gatheir Miller, Yung Sung Cheng, Richard J. Traub, Tom T. Little, and Raymond A. Guilmetes 34.			
Mary Ann Parkhurst and Raymond Aerosol Sampling System for Depleted Uranium Particles is Environment Thomas D. Holmes, Raymond A.	Collection of Capstone n a High-Energy Guilmette, Yung Sung Cheng,	207	Capstone Depleted Uranium Aerosol Blokinetics, Concentrations, and Doses Raymond A. Goilnette, Gubrie Miller, and Mary Ann Parkhurst Assessing the Renal Toxicity of Capstone Depleted Uranium Oxides and Other Uranium	3		
Mary Ann Parkhurst, and Mark D. Calculating Capstone Depleted Concentrations from Beta Acti Frances Szrom, Gerald A. Falo, Jeffrey J. Whicker, and David P.	Uranium Aerosol ivity Measurements Mary Ann Parkhurst,	238	Compounds Laurie E. Rozell, Pletcher F. Hahn, Robyn E. Lee, and Mary Ann Parkhunst Radiological Risk Assessment of Capatone Depleted Uranium Aerosols	3		
Physicochemical Characterizat Uranium Aerosols I: Uranium a Function of Time and Partic Mary Ann Parkhurst, Yung Sung (Richard J. Traub	Concentration in Aerosols as le Size	251	Fletcher F. Hahn, Laurie E. Roszell, Eric G. Daxon, Ray A. Guilmette, and Mary Ann Parkhurst Inhalation and Ingestion Intakes with Associated Dose Estimates for Level II and Level III Personnel Using Capstone Study Data	3		
Physicochemical Characteriza Uranium Aerosols II: Particle Function of Time Yung Sung Cheng, Judson L. Ken and Mary Ann Parkhurst	e Size Distributions as a	266	Frances Szrom, Gerald A. Falo, Gordon M. Lodde, Mary Ann Parkhunst, and Eric G. Daxon Application of Capstone Depleted Uranium Aerosol Risk Data to Military Combat Management Eric G. Daxon, Mary Ann Parkhunst, Mark A. Melanson,	3		
Physicochemical Characteriza Uranium Aerosols III: Morph Analyses Kenneth M. Krupka, Mary Ann Pa	hologic and Chemical Oxide arkhurst, Kenneth Gold,		and Laurie E. Roszell Conclusions of the Capstone Depleted Uranium Aerosol Characterization and Risk Assessment Study May Ann Parkhunt and Raymond A. Goilmette	3		
Brace W. Arey, Evan D. Jenson, a Physicochemical Characteriza Uranium Aerosols IV: In Vita	ation of Capstone Depleted ro Solubility Analysis	276	OTHER CONTENT			
Raymond A. Guilmette and Yung	Sung Cheng	292	News and Notices	X		

UNCLASSIFIED





LESSONS LEARNED CAPSTONE DU STUDY AND HUMAN HEALTH RISK ASSESSMENT

Information from the Capstone DU reports applied almost daily to current military operations regarding occupational and environmental health issues

- Army Training Ranges
- Historical weapon systems with DU
- Southwest Asia military operations
- Radioactive waste clean-up and disposal
- BRAC installations and environmental issues
- Soldier Post-Deployment Health Assessments
- DU Activist challenges





OVERVIEW DEPLETED URANIUM (DU) AND ITS HEALTH EFFECTS (I)

- An overview of uranium health effects may be found in the US Agency for Toxic Substances and Disease Registry (ATSDR) Toxicological Profile for Uranium (see http://www.atsdr.cdc.gov/toxprofiles/tp150.html, a 1999 update to the original profile published in May 1989)
- Quote: "While natural and depleted uranium are considered chemically toxic, they are not considered a [significant] radiation hazard"
- Discussions are currently ongoing among several Federal Agencies to fund ATSDR to produce an updated addendum to this Uranium Toxicological Profile





OVERVIEW DEPLETED URANIUM (DU) AND ITS HEALTH EFFECTS (II)

- The International Agency for Research on Cancer (World Health Organization) Monographs, Volume 78, Ionizing Radiation, and Volume 78, Part 2: Some Internally Deposited Radionuclides (2001), are another basic source of information on radiological effects
- See http://monographs.iarc.fr/ENG/Monographs/vol78/mono78.pdf
- M. A. McDiarmid, et. al., Surveillance Results of Depleted Uranium –Exposed Gulf War I Veterans: Sixteen Years of Follow-Up, Journal of Toxicology and Environmental Health, Part A, 72: 14–29, 2009





OVERVIEW DEPLETED URANIUM (DU) AND ITS HEALTH EFFECTS (III)

- "As part of a longitudinal surveillance program, 35 members of a larger cohort of 77 Gulf War I veterans who were victims of depleted uranium (DU) "friendly fire" during combat underwent a 3-day clinical assessment at the Baltimore Veterans Administration Medical Center (VAMC)"
- "Sixteen years after first exposure, this cohort continues to excrete elevated concentrations of urine U as a function of DU shrapnel burden. Although subtle trends emerge in renal proximal tubular function and bone formation, the cohort exhibits few clinically significant U-related health effects."





OVERVIEW DEPLETED URANIUM (DU) AND ITS HEALTH EFFECTS (IV)

- The National Academy of Sciences Institute of Medicine recently updated its literature review on the health effects of DU and other Gulf War environmental hazards:
 - "Gulf War and Health Volume 8 Update of Health Effects of Serving in the Gulf War" (IOM, NAP, April 2010)"
 - In this report, the IOM determines that DU is not a significant source of occupational exposure
- •The Department of Veterans Affairs (VA) published a notice in the Federal Register (FRN) in March 2010 on its decision related to a 2006 Institute of Medicine (IOM) report on the potential health effects of DU (IOM, NAP, Report, Gulf War and Health, Volume 4: Health Effects of Serving in the Gulf War, 2006) and the Volume 8 Update






OVERVIEW DEPLETED URANIUM (DU) AND ITS HEALTH EFFECTS (V)

- The VA's FRN states that the IOM also concluded there
 was limited or suggestive evidence of no association
 between uranium and clinically significant renal
 dysfunction and between uranium and lung cancer at
 specified cumulative internal doses
- Therefore, the VA has decided NOT to service-connect any diseases related to DU, as listed in the IOM report
- See Federal Register/Vol. 75, No. 45 /VA /Notice , 9 March 2010
- http://edocket.access.gpo.gov/2010/pdf/2010-4882.pdf





SCREENING BIOASSAY DATA QUALITY OBJECTIVES



PROGRAM GOAL

Identify all personnel with elevated levels of DU and embedded fragments that contain DU (and tungsten composites) for referral to the VA Medical DU Follow-Up Program

- Laboratory performs urine uranium analysis using Inductively Coupled Plasma-Mass Spectroscopy (ICP-MS) on MTF urine specimens
- Laboratory determines isotopic ratios for DU in urine specimen at detection limits as low as possible with laboratory method
- Health Physics Program reviews laboratory results and prepares a health risk assessment report for requesting healthcare provider at MTF
- AMEDD, through the DoD Deployment Health Clinical Center, refers patients with elevated urine uranium levels indicative of DU to VA Medical DU Follow-Up Program





AMEDD DU POLICY

Policy for DU Bioassay and Fragment Analysis:

- Memorandum, U.S. Army Medical Command, MCPO-SA, OTSG/MEDCOM Policy Memo 09-038, 18 June 2009, subject: Medical Management of Army Personnel Exposed to Depleted Uranium (DU).
- Supersedes OTSG/MEDCOM Policy 07-022, 26 June 2007).
- See http://www.pdhealth.mil/du.asp#army





OIF DU Results Summary (as of 30 Sep 10)

- Urine bioassay specimens
 - Over 2350 individuals screened
 - Over 2750 analyses
 - 8 individuals with DU in specimen
 - 5 elevated above NHANES, 3 in NHANES range
 - Preliminary dose estimates for 6 Soldiers
 - E(50) range: 0.00001Sv (0.001 rem) to 0.012 Sv (1.2 rem)
 - U kidney concentration: 0.003 to 11.1 μg U per gram kidney tissue
 - 4 Soldiers referred to VA DU Follow-Up Program
- Fragment analyses
 - 80 fragments from 69 individuals
 - 3 fragments from 2 individuals positive for U
- Urine uranium results consistent w/ CDC 3rd NHANES data/higher than 4th NHANES data










Comparison of Uranium Specimen Values to 4th NHANES Data and Laboratory Screening Level (as of 30 September 2010)







New Metal Fragment Policy

- Memorandum, Assistant Secretary of Defense for Health Affairs, HA Policy 07-029, 18 December 2007, subject: Policy on Analysis of Metal Fragments Removed from Department of Defense Personnel
- Memorandum, U.S. Army Medical Command, MCPO-SA, OTSG/MEDCOM Policy Memo 10-041, 29 June 2010, subject: Management of Metal Fragments Removed from Army Personnel
- Application of lessons learned from DU Policy experiences
 - Eventual integration of two policies into a Heavy Metals Policy
 - Eventual electronic patient records





Metal Fragment Policy

- Specifically, as part of the new DOD and Army MEDCOM embedded metal fragment policies, USAIPH received 235 embedded metal fragments from 147 patients from 147 October 2009 through 30 September 2010
- Cumulative analysis results (under the embedded metal fragment policies) are provided in the following table and graph





Summary of Laboratory Results Removed Embedded Metal Fragments Cumulative Reporting Period: 18 September 2008 – 30 September 2010


Report Period		# fragments received	# fragments analyzed*	# packages sent out ¹	Content: common metals ²	Content: lead ³	Content: any tungsten ⁴	Content: tungsten- nickel- cobalt	Content: High content Nickel
18 Sep 08 through 30 Sep 10	147	235	207	137	182	15	2	0	0

Notes:

- * One fragment not analyzed due to fragment size and composition
- 1. "Packages Sent" listed by # of fragments, but actually are combined per person when sent.
- 2. "Common Metals" are considered anything other than lead, tungsten or high content nickel. There are no separate packages (with Cover Memo interpretation and fact sheet) to separate out these metals for special consideration
- 3. Thirteen (13) fragments with primary content lead. There are 2 more with small to trace amounts of lead on common metals.
- 4. "Any Tungsten" often includes surface tungsten only. Tungsten content analyzed through 30 September 2010 has been considered minor and is counted under "Common Metals."











Embedded Metal Fragment Assessment

For the 207+ fragments analyzed since implementation of HA Policy 07-029:

- None are depleted uranium (several fragments submitted previously were identified as DU)
- None are radioactive
- Tungsten has been detected, but in relatively low concentrations; primary tungsten alloy (composite) of concern has not been identified





PROPOSED DISCUSSION TOPICS

DU researchers, medical planners, and combat operations staff are suggesting follow-up:

- Additional follow-up studies of exposed populations have the
 potential to improve knowledge of the health effects of DU. To permit
 an adequate assessment of the risks of cancer, renal toxicity, and
 other possible health effects faced by DU-exposed soldiers, a careful
 follow-up of the exposed groups should be continued, including the
 cohort now being followed by the Department of Veterans Affairs.
- An examination of subgroups of Soldiers with high, medium, and low exposure to uranium, with appropriate matching on other risk factors, should be implemented for selected health-related end points and biomarkers.
- The DU researchers should ask the military medical planners and combat operations staff what information is still needed to protect the Soldier and other Servicemembers on the battlefield.





References

- Memorandum, Assistant Secretary of Defense for Health Affairs, HA Policy 03-012, 30 May 2003, subject: Policy for the Operation Iraqi Freedom Depleted Uranium (DU) Medical Management. (www.ha.osd.mil/policies/2003/03-012.pdf)
- Memorandum, U.S. Army Medical Command, MCPO-SA, OTSG/MEDCOM Policy Memo 09-038, 18 June 2009, subject: Medical Management of Army Personnel Exposed to Depleted Uranium (DU). (Supersedes OTSG/MEDCOM Policy 07-022, 26 June 2007). (http://www.pdhealth.mil/du.asp#army)
- Centers for Disease Control and Prevention, National Center for Environmental Health, Fourth National Report on Human Exposure to Environmental Chemicals, National Health and Nutrition Examination Surveys (NHANES), December 2009 (www.cdc.gov/exposurereport)
- DU Questionnaire: DD Form 2782 Test, Feb 2004 and DD Form 2782-1 Test, Feb 2004. Available at: www.pdhealth.mil/du.asp
- DU information available at: http://fhp.osd.mil/du/
- National Council on Radiation Protection and Measurements (NCRP)
 Report 156, Development of a Biokinetic Model for RadionuclideContaminated Wounds for Their Assessment, Dosimetry and
 Treatment, 2006. (http://www.ncrponline.org/Publications/156press.html)



SOLDIER EXPOSURE TO DEPLETED URANIUM AND HUMAN HEALTH RISK ASSESSMENT METHODOLOGY UPDATE - 2010

DAVID P. ALBERTH
SENIOR HEALTH PHYSICIST AND MASTER CONSULTANT

FRANCES SZROM, CHP

HEALTH PHYSICIST AND MASTER CONSULTANT

US ARMY INSTITUTE OF PUBLIC HEALTH
DEPLETED URANIUM (DU) HEALTH EFFECTS AND POLICY WORKSHOP

AFRRI, Bethesda, Maryland
4 November 2010



UNCLASSIFIED